

Unusual condensates in quark and atomic systems*

B.Kerbikov,[†]

State Research Center

Institute of Theoretical and Experimental Physics,
Moscow, 117218, Russia

Abstract

In these lectures we discuss condensates which are formed in quark matter when it is squeezed and in a gas of fermionic atoms when it is cooled. The behavior of these two seemingly very different systems reveals striking similarities. In particular, in both systems the Bose-Einstein condensate to Bardeen–Cooper–Schrieffer (BEC-BCS) crossover takes place.

1 Introduction

In these lectures I will review the striking discoveries of the new phases of the quark and atomic matter. Squeezing the quark matter and cooling the gas of fermionic atoms lead to the formation of condensates. Condensates in both systems exhibit similar properties. The most interesting point is the crossover between different regimes of condensation, which takes place under the variation of certain external parameters.

At first sight the dense quark matter and cold fermionic atoms have nothing in common as one might conclude from Table 1.

*Lectures delivered at 8th Moscow school of Physics (33rd ITEP Winter School of Physics)

[†]E-mail:borisk@itep.ru

Table 1Comparison of density and temperature in quark and atomic systems¹

	density	temperature
quarks	$> 10^{39} cm^{-3}$	$\lesssim 5 \cdot 10^{11} K$
fermionic atoms	$(10^{13} - 10^{14}) cm^{-3}$	$\sim (10^{-7} - 10^{-8}) K$

The numbers presented in Table 1 will be discussed below, but some comments are in order already at this point. The quark density $n \sim 10^{39} cm^{-3}$ is not high, it is only three times larger than the density of quarks in normal nuclear matter. When the density reaches this value the gap equation develops a nontrivial solution, which was commonly interpreted as the onset of the BCS-like phase. We shall argue that this is a misleading conclusion and the BCS regime corresponds to much higher densities.

The density of fermionic atoms $n \sim (10^{13} - 10^{14}) cm^{-3}$ is very low, it is about five orders of magnitude lower than the density of the air. Such a low density is needed in order to suppress the formation of molecules and clusters. To obtain quantum condensation in such a dilute system it should be cooled to extremely low temperatures $\sim (10^{-7} - 10^{-8}) K$ since roughly speaking the temperature of BEC scales as $n^{2/3}$. The fascinating aspect about ultracold fermionic atoms is that the strength of the interatomic interaction can be tuned using the magnetic Feshbach resonance (see below).

Intense discussion of the unusual properties of quark matter at finite density (color superconductivity) has started in 1998 after the publication of the pioneering papers [1, 2]. Condensates in a gas of ultracold fermionic atoms have been experimentally created in 2003 [3, 4, 5]. The idea that the BCS regime in quark system sets in as a result of the BEC-BCS crossover was formulated in 2001 [6, 7]. Hundreds of papers have been published in recent years on quark systems at nonzero density and on ultracold fermionic atoms. The reader willing to get deeper and broader knowledge on both subjects may address review papers [8, 9].

¹We remind that 1 eV = 11606 K

2 Two types of condensation: BEC and BCS

It is not easy to give a universal definition of a condensate, though in any concrete situation this notion is well understood. The key properties of a condensate are:

- (a) Long-range order.
- (b) Macroscopic quantum coherence, i.e., the possibility to regard the condensate as a single quantum entity described by the macroscopic wave function.

The type of condensate (the character of quantum degeneracy) depends upon the spin of the particles. In a system of bosons BEC takes place when de Broglie thermal wavelength $\lambda_B = (2\pi/mT)^{1/2}$ is of the order of interparticle distance, $\lambda_B n^{1/3} \sim 1$. More accurate analysis yields the degeneracy condition in the form $\lambda_B^3 n = s\zeta(3/2)$, or

$$T_c(\text{BEC}) = \frac{2\pi}{m} \left[\frac{n}{s\zeta\left(\frac{3}{2}\right)} \right]^{2/3}. \quad (1)$$

Here s is the number of possible spin states, $\zeta(3/2) \simeq 2.6$ is the Riemann zeta function, and we set $k_B = \hbar = 1$.

BEC has a very long history. The best known example of this regime is ^4He below the so-called λ -point at 2.17 K. Condensations of π - and K -mesons were discussed as possible states of neutron stars. BEC-like behavior is present in some scenarios of high T_c superconductivity. Ten years ago BEC was observed in trapped gases of bosonic alkali atoms ^{87}Rb and ^{23}Na [10]-[12]. In these lectures we shall discuss BEC and BCS condensation in the systems of fermionic atoms and quarks. A natural question at this point would be : "How is it possible to talk about BEC for fermions?" The answer to this question is the following. Interaction between fermions may lead to pairing and the resulting composite bosons may undergo BEC – "A molecular Bose–Einstein condensate emerges from a Fermi sea" [3]. This was realized experimentally for two fermionic alkalies – ^6Li and ^{40}K [3, 4, 5]. Both have even number of nucleons and one electron outside the closed shell and therefore they are fermions. For $^6\text{Li}_2$ molecules with the density $n = 10^{13} \text{ cm}^{-3}$ Eq.(1) gives $T_c \simeq 5 \cdot 10^{-7} K$ in a fair agreement with

the experimental result $T_c = 2.8 \cdot 10^{-7} K$ [4]. In reality, however, (1) is not applicable to the trapped gases. Condensation in this case means macroscopic population of the lowest trap state and the critical temperature is given by $T_c = \omega (N/\zeta(3))^{1/3} \simeq 0.94 \omega N^{1/3}$ [13], where N is the number of particles in the trap (typically $N \sim 10^5$), ω is the average oscillation frequency of the trap ($\omega \sim \text{few } nK$), $\zeta(3) \simeq 1.2$. Thermodynamic limit corresponds to $N \rightarrow \infty, \omega \rightarrow \infty, N\omega^3 = \text{const}$ [13]. Other important factors making the thermodynamic of the trapped gas nontrivial are the trap geometry, inhomogeneity of the system, and the effects of the interaction between the molecules [13]-[16].

The observation of BEC for fermionic atoms is not the main reason why the experiments [3, 4, 5] generated enormous excitement. The striking discovery was the creation of a state with another type of pairing between the atoms, namely the BCS condensate and the observation of the crossover between BEC and BCS regimes [3-5,9]. Before turning to the crossover phenomena let us remind few basic facts about Cooper pairing and BCS.

Cooper pairing is the cornerstone of the theory of superconductivity – one of the most elegant theories created in the last century. The nature of Cooper pairing is different from that of molecular-like states discussed above². The size of the Cooper pair is characterized by the coherence length ξ which is macroscopic, i.e. much larger than the interatomic distance, $\xi \sim 10^{-4}$ cm, while $a \sim 10^{-8}$ cm. The important dimensionless parameter is $n^{1/3}\xi$, where n is the density of fermions. For a typical superconductor the density of electrons is $n \sim 10^{22} \text{ cm}^{-3}$, so that $n^{1/3}\xi \gtrsim 10^3$. Alternatively, instead of $n^{1/3}$ we may use the Fermi momentum $k_F = (3\pi^3 n)^{1/3}$. Thus the BCS regime is characterized by the dimensionless parameter

$$k_F \xi \gtrsim 10^3. \quad (2)$$

Equation (2) means that Cooper pairs strongly overlap with each other and the problem is essentially many-body. In coordinate space the wave function of the Cooper pair is proportional to $(\sin k_F r / k_F r) \exp(-r/\xi)$ and therefore it has $\gtrsim 10^3$ nodes. We see that pairing in the BCS regime is very different from the usual molecular-like one. The dynamical mechanism leading to the Cooper phenomenon includes three important elements:

²Compact electron pairs were introduced [17] in an attempt to explain superconductivity couple of years before the BCS theory was formulated. Sometimes compact molecular-like states are called Schafroth pairs.

- (i) Attraction between particles.
- (ii) Existence of the the Fermi surface.
- (iii) Interaction must be concentrated within a thin layer of momentum space around the Fermi surface.

Attraction between electrons in a superconductor is due to the interaction with the lattice (phonon mechanism). The characteristic energy scale of this interaction is the Debye frequency $\omega_D \simeq \frac{1}{2} (m/M)^{1/2} Ry \sim 10^{-2}$ eV, where m and M are the electron and ion, masses³. On the other hand, the typical Fermi energy is $E_F = k_F^2/2m \simeq (3\pi^2 n)^{2/3}/2m \simeq 2$ eV. Thus $\omega_D \ll E_F$ and we see that the interaction in the BCS regime is really concentrated within a thin layer around the Fermi surface. Electrons interact in two dimensions instead of three, the sum $\sum_k \mathbf{k}^{-2}$ logarithmically diverges at small momenta leading the Cooper instability and to the gap in the spectrum.

At this point we wish to demonstrate the importance of the point (iii), or the hierarchy $\omega_D \ll E_F$, in a more formal way. Use will be made of the technique standard for the quantum field theory. Similar language was used, e.g., in Refs. [19, 20]. We start with the partition function in the form of the path integral over the fermion fields

$$Z = \text{tr } e^{-\beta H} = \int D\psi^+ D\psi \times \\ \times \exp \left\{ - \int_0^\beta d\tau \int d\mathbf{r} \left[\sum_\sigma \psi_\sigma^+ \left(\frac{\partial}{\partial \tau} - \frac{\nabla^2}{2m} - \mu \right) \psi_\sigma - g \psi_\sigma^+ \psi_{-\sigma}^+ \psi_{-\sigma} \psi_\sigma \right] \right\}. \quad (3)$$

Here σ is the spin variable, $\mu \simeq E_F$ due to (iii)⁴. The constant g has the dimension of m^{-2} , the dimension of $\psi_\sigma(\mathbf{r}, \tau)$ is $m^{3/2}$. From now on we shall consider stationary homogeneous superconductor and assume that the four-fermion interaction is local. The last term in the exponent in (3) is quartic in the fermion fields making the direct integration impossible. This difficulty can be overcome by Hubbard–Stratonovich trick (bosonization). To decouple the quartic interaction one writes

$$\exp \left\{ g \int_0^\beta d\tau \int d\mathbf{r} \psi_\sigma^+ \psi_{-\sigma}^+ \psi_{-\sigma} \psi_\sigma \right\} =$$

³Close to the Fermi surface this attraction overcomes Coulomb repulsion – see, e.g. [18].

⁴We stress that $\mu \simeq E_F$ is true only in the BCS regime.

$$= \int Dd^* Dd \exp \left\{ - \int_0^\beta d\tau \int d\mathbf{r} \left(\frac{|d|^2}{g} - d\psi_\sigma^+ \psi_{-\sigma}^+ - d^* \psi_{-\sigma} \psi_\sigma \right) \right\}. \quad (4)$$

Here we have introduced complex scalar field d . The quantity $|d|$ will acquire the meaning of the superconducting order parameter. From the Lagrange equation of motion for d^* we see that $d = g\psi_{-\sigma}\psi_\sigma$, so that d has a dimension of mass as it should be for the gap. Before we insert (4) into (3) and proceed on with our derivation of Z , some important remarks should be made.

The initial microscopic Hamiltonian in (3) is proportional to $\psi_\sigma^+ \psi_{-\sigma}^+ \psi_{-\sigma} \psi_\sigma$. According to Wick's theorem it has to be factorized as $\langle \psi_\sigma^+ \psi_{-\sigma} \rangle \langle \psi_{-\sigma}^+ \psi_\sigma \rangle$. In such a form the Hamiltonian conserves the number of particles and does not lead to superconductivity. Equation (4) corresponds to an alternative factorization $\langle \psi_\sigma^+ \psi_{-\sigma}^+ \rangle \langle \psi_{-\sigma} \psi_\sigma \rangle$. In a normal, non-condensed state such anomalous averages correspond to the off-diagonal long range order (ODLRO) – a concept introduced by Yang [21], and relevant both for BEC and BCS regimes. The state with ODLRO is characterized by a long range correlation

$$S(\mathbf{r}_1, \mathbf{r}_2) \propto d^*(\mathbf{r}_1)d(\mathbf{r}_2), \quad |\mathbf{r}_1 - \mathbf{r}_2| \rightarrow \infty. \quad (5)$$

The wave function in ODLRO regime is a coherent state with fluctuating number of particles [22, 23]

$$|\Psi\rangle = \exp \left\{ -\frac{N}{2} + \sqrt{N}b^+ \right\} |0\rangle, \quad (6)$$

where b^+ is a pair creation operator. For a metallic superconductor $N \sim 10^{20}$ and the uncertainty in the number of particles is $\Delta N/N \sim \sqrt{N}/N \sim 10^{-10}$ which is not important. However, for the condensate of fermionic atoms $N \sim 10^5$, $\Delta N/N \gtrsim 10^{-3}$ and this might be physically significant. An interesting open question is the role of phase/particle number uncertainty relation $(\Delta\theta)(\Delta N) \gtrsim 1/2$ [24] for condensate with rather low value of N . Another open question is the phase variation in BEC-BCS crossover.

Now we return to the evaluation of the partition function. We substitute (4) into (3) and rewrite the result in the matrix Nambu-Gorkov representation.

$$Z = \int D\psi^+ D\psi Dd^* Dd \exp \left\{ - \int_0^\beta d\tau \int d\mathbf{r} \left[\frac{|d|^2}{g} + (\psi_\sigma^+, \psi_{-\sigma}) \begin{pmatrix} R & d \\ d^* & R^T \end{pmatrix} \begin{pmatrix} \psi_\sigma \\ \psi_{-\sigma}^+ \end{pmatrix} \right] \right\}. \quad (7)$$

Here $R = \left(\frac{\partial}{\partial \tau} - \frac{\nabla^2}{2m} - \mu \right)$, $R^T = \left(\frac{\partial}{\partial \tau} + \frac{\nabla^2}{2m} + \mu \right)$. Equation (7) is simply a mathematical construct completely equivalent to (3)-(4). In order to follow back the route from (7) to (3)-(4) one should keep in mind the anticommutativity of fermion fields and be ready to integrate by parts the operator R^T . Next we integrate over the fermion fields using the fact that (7) contains a quadratic form in the exponent and the integral is of a Gaussian type. The result reads

$$Z = \int Dd^* Dd \exp \left\{ -\frac{1}{g} \int_0^\beta d\tau \int d\mathbf{r} |d|^2 - \beta \Omega_B(T, \mu, d, d^*) \right\}, \quad (8)$$

where Ω_B is the Bogolubov functional

$$\beta \Omega_B = -\text{tr} \ln \begin{pmatrix} R & d \\ d^* & R^T \end{pmatrix} = - \int_0^\beta d\tau \int d\mathbf{r} \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{1}{\beta} \sum_{\omega_n} \ln(\omega_n^2 + |d|^2 + E^2(k)). \quad (9)$$

Here summation runs over the fermionic Matsubara modes $\omega_n = \frac{\pi}{\beta}(2n+1)$ [25], and $E(k) = \frac{k^2}{2m} - \mu$.

The value of the gap d is given by the extremum condition for the complete action in the exponent of (8)

$$d = -\frac{g}{V_3} \frac{\delta \Omega_B}{\delta d^*}, \quad (10)$$

where $V_3 = \int d\mathbf{r}$. The Matsubara summation in $\delta \Omega_B / \delta d^*$ is performed using the relation

$$T \sum_{\omega_n} \frac{2\varepsilon(k)}{\omega_n^2 + \varepsilon^2(k)} = th \frac{\varepsilon(k)}{2T}. \quad (11)$$

Here $T = 1/\beta$, $\varepsilon^2(k) = E^2(k) + |d|^2$. From (9-11) we obtain the self-consistency relation

$$1 = \frac{g}{2} \int \frac{d\mathbf{k}}{(2\pi)^3} \varepsilon^{-1}(k) th \frac{\varepsilon(k)}{2T}. \quad (12)$$

At this point the BCS condition (iii) formulated above comes into play. Attractive force between electrons due to phonons acts within a thin layer around the Fermi surface $-\omega_D \leq E \leq \omega_D$. Therefore we may cut the integral in (12) and write

$$\int \frac{d\mathbf{k}}{(2\pi)^3} \simeq \frac{mk_F}{2\pi^2} \int_{-\omega_D}^{\omega_D} dE = 2N(0) \int_0^{\omega_D} dE, \quad (13)$$

where $N(0) = mk_F/2\pi^2$ is the density of states on the Fermi surface. Then we substitute (13) into (12), take the limit $T \rightarrow 0$ and obtain the famous BCS solution

$$d = 2\omega_D \exp\left(-\frac{1}{gN(0)}\right) = 2\omega_D \exp\left(-\frac{\pi}{2k_F|a|}\right), \quad (14)$$

where the scattering length is introduced according to $g = 4\pi|a|/m$ ⁵. The BCS regime corresponds to $a < 0$ if we use the standard sign convention $k \cot \delta \rightarrow -a^{-1}$. We can estimate $|a|$ for a typical superconductor with $\omega_D \sim 10^2 K$ and $d \sim T_c \sim \text{few } K$. Then according to (14) $gN(0) \simeq 0.3$ and $|a| \sim 10^{-8} \text{ cm} \sim k_F^{-1}$.

3 Condensation in Quark Systems and in Atomic Fermi Gases

With BEC and BCS ideas and technique at hand we may try to understand to what extent are they applicable to the systems of quarks and fermionic atoms.

Consider the system of u - and d -quarks with density (3-5) times larger than the density of quarks in normal nuclear matter (see Table 1). The corresponding value of the chemical potential is around $\mu \simeq 0.4 \text{ GeV}$ [28]. Such densities are typical for neutron stars. Strange quark starts to participate in pairing at much higher densities corresponding to $\mu \gg m_s \simeq 150 \text{ MeV}$. Quark matter with such moderate density got the name of 2 SC phase, where 2 stands for the number of flavors and SC is a shorthand form of "superconducting color". In enormous number of publications one finds the statement that the BCS regime really sets in starting from $\mu \simeq 0.4 \text{ GeV}$. This conclusion relies on the fact that at such values of μ the gap equation of the type (10) acquires a nontrivial solution (see [28] and references in [8]). From the preceding discussion it should be clear, however, that a nonzero value of the gap is only a signal of the presence of fermion pairs. Depending on the dynamics of the system, on the fermion density, and on the temperature, such pairs may be either stable, or fluctuating in time, may form a BCS condensate or dilute Bose gas, or undergo a Bose condensation. We

⁵The rigorous definition of the scattering length for the contact interaction entering into (3) may be found in [26, 27].

have seen that the BCS regime requires for its realization rather special conditions (i)-(ii)-(iii) formulated above. Let us show that these requirements are not met in the 2SC phase [6, 7].

Scalar diquark in the 2SC regime is a would-be Cooper pair. We recall that the spin structure of the order parameter in normal superconductor has the form $d \sim \varphi_\sigma(1)(i\sigma_2)_{\sigma\sigma'}\varphi_{\sigma'}(2)$. Similarly for scalar diquark in $\bar{3}$ color state we may write [8] $d_\alpha \sim \delta_{\alpha 3}\bar{q}_{\beta i}^c \varepsilon_{\beta\gamma 3}(\tau_2)_{ij}\gamma_5 q_{\gamma j}$, where the Greek indices stand for color, Latin ones for flavor, $q^c = Cq^T$, C is the charge conjugation operator, and the presence of γ_5 makes the diquark a scalar (like σ_2 for electron pair). Diquark has a certain orientation in color space (e.g., along the third axis $\delta_{\alpha 3}$), therefore $SU(3)_c$ gauge symmetry is broken, five of eight gluons are massive. On the contrary, chiral symmetry is restored. The symmetry pattern changes in the ultra-high density region when the strange quark enters the game. In the $\mu \rightarrow \infty$ limit the real BCS regime sets in (though with some peculiarities [8]). We left out this topic since our main interest is focused on the BCS-BEC interplay at moderate density.

The dynamics of quark matter at moderate density is characterized by the typical momenta $k \sim \mu \lesssim 1$ GeV and by typical distances $r \sim \mu^{-1} \sim 1$ fm. In this domain theory faces the well known difficulties of nonperturbative QCD and use is made of models like NJL or instanton gas. The form of the Bogolubov functional is practically independent of the model and reads [29]

$$\begin{aligned} \beta\Omega_B &= -\frac{1}{2}\text{tr} \ln \begin{pmatrix} i\partial_\mu \gamma_\mu - i\mu\gamma_4 & d\hat{O} \\ d^*\hat{O}^+ & -i\partial_\mu^T \gamma_\mu^T + i\mu\gamma_4 \end{pmatrix} = \\ &= -\text{tr} \ln(-\hat{k} - i\mu\gamma_4) - \frac{1}{2}\text{tr} \ln \left\{ 1 + \frac{\hat{O}\hat{O}}{\hat{k}_+\hat{k}_-} |d|^2 \right\}, \end{aligned} \quad (15)$$

where $\hat{k} = k_\mu \gamma_\mu$, $\hat{O} = \varepsilon_{\beta\gamma 3}(\tau_2)_{ij} C \gamma_5$, $k_\pm = (\mathbf{k}, k_4 \pm i\mu)$.

The self-consistency relation (see(12)) at $T = 0$ has the form (for zero current quark masses)

$$1 = \frac{g}{2} \int \frac{d\mathbf{k}}{(2\pi)^3} \{ \varepsilon_+^{-1}(k) + \varepsilon_-^{-1}(k) \}, \quad (16)$$

where $\varepsilon_\pm^2(k) = (k \mp \mu)^2 + |d|^2$.

Acting essentially within the same scheme which led from (9) to (14) one obtains the result $d \simeq 0.1$ GeV almost independently of the model [1, 2, 8, 28]. The roles of $N(0)$, ω_D and E_F are played by $N(0) = 2\mu^2/\pi^2$, momentum

cutoff $\Lambda \simeq 0.7$ GeV, and $E_F \simeq \mu \simeq 0.5$ GeV. Let us compare these numbers with what we have in the standard BCS picture

$$BCS \quad d : \omega_D : E_F \simeq 1 : 10^2 : 10^4 \quad (17)$$

$$2SC \quad d : \Lambda : \mu \simeq 1 : 8 : 5. \quad (18)$$

We conclude that the hierarchy of scales inherent in the BCS regime is badly broken in the 2SC phase. As we have seen there is another important parameter in the BCS scenario, namely (see(2)) $k_F \xi \gtrsim 10^3$. Let us estimate it for the 2SC regime. Table 1 shows that $n^{1/3} \simeq 0.2$ GeV in the 2SC phase⁶. As for the value of ξ , we may only rely on some estimates since rigorous calculations are hardly possible in the nonperturbative QCD region. One should also keep in mind distinction between the correlation length and the pair size. The two quantities coincide in the BCS regime [30], while in BEC region the pair size is smaller than the coherence length [31, 32]. The naive estimate of ξ may be done as follows. The energy spread of the correlated pair of quarks is $\delta E \sim d$, quarks are relativistic, hence $\delta p \sim d$, and therefore $\xi \sim 1/d \sim 2$ fm. Using the Klein-Gordon equation for the quark pair [33] one can obtain a better estimate $\xi \simeq (\sqrt{3}d)^{-1} \lesssim 1$ fm⁷. Therefore we have ($k_F \simeq \mu$)

$$k_F \xi \simeq 2. \quad (19)$$

Comparing with (2) we see that this value is about three orders of magnitude less than in the BCS regime. The result (19) indicates that the 2SC phase is in halfway between the BEC and BCS regions. The role of fluctuations in the crossover domain is very important [34]. The significance of fluctuations is directly seen from the estimate of the Ginzburg–Levanyuk parameter

$$Gi \sim (k_F \xi)^{-4} \sim 1/16. \quad (20)$$

For the ordinary superconductor $Gi \sim 10^{-12} - 10^{-14}$. Two kinds of fluctuations should be considered – that of the order parameter d , and that of the gluon field A [35]. The dominant ones are the gluon field fluctuations since their correlation length T_g is about 0.2 fm [36], i.e., about 5 times

⁶This value is obtained according to $n = -\partial\Omega_B/\partial\mu$, but it is close to the result for free degenerate quarks $n = N_c N_f \mu^3/3\pi^2$.

⁷Recall that in the BCS theory $\xi \simeq v_F/\pi d$ [23].

smaller than the correlation length ξ for d . Fluctuation diamagnetism [34] gives rise to the substantial shift of the critical temperature [37, 38, 39]

$$T'_c = T_c(1 - g^2 \xi^2 \langle A^2 \rangle), \quad (21)$$

where g is the strong coupling constant, A – the gluon field, $\langle \dots \rangle$ is the average over the quark fields, $T'_c \simeq 0.75T_c$.

It is clear that further work beyond the Mean Field Approximation is needed before the precise dynamics of the 2SC phase is finally elucidated. In this respect very helpful might be the rapid progress taking place now in the study of similar problems for ultracold fermionic atoms. The central point here is the experimental possibility to tune the interatomic scattering length via the magnetic Feshbach resonance [40, 41]. The scattering length entering into equation (14) may change its value and sign as a function of the external magnetic field. Near the resonance the dependence of the scattering length on the magnetic field reads

$$a(B) = a_0 \left(1 - \frac{\Delta}{B - B_r} \right), \quad (22)$$

where a_0 is the “background” scattering length, Δ is some constant, and B_r is the value of the magnetic field corresponding to the resonance at threshold ($a(B_r) \rightarrow \infty$). The behavior of the scattering length as a function of the interaction strength g is well known from quantum mechanics. With g increasing the scattering length first tends to $a \rightarrow -\infty$, then changes to $a \rightarrow +\infty$ (i.e., pass through a resonance at threshold) and then tends to $a^{-1} \rightarrow \infty$ in the strong coupling limit when the scattering length approximation is no longer legitimate. In nuclear physics such an evolution may be regarded as a transition from the np virtual state with antiparallel spins and $a \simeq -24$ fm to the deuteron with $a \simeq 5.4$ fm and $\varepsilon_b \simeq 2.2$ MeV. In physics of cold fermion atoms the region $a < 0$ is called “BCS side of resonance” and the region $a > 0$ got the name “BEC side of resonance”. We also know that in the scattering length approximation the binding energy is equal to $\varepsilon_b = (ma^2)^{-1}$ with m being the mass of each constituent forming the pair. At this point the “connection problem” naturally arises: is it possible to connect by a certain continuous transformation the BCS solution (14) and the scattering length solution presented above? Before we formulate a positive answer to this question a remark concerning manipulations with the scattering length approximation and with the two-body interaction is general

should be made. The BCS solution (14) is of essentially many-body nature. If we consider an isolated pair of electrons near the Fermi surface and solve the eigenvalue problem in weak coupling limit and in thin layer approximation, we would obtain the result similar to (14) but with twice as large factor in the exponent. The overlap of many Cooper pairs is important. The solution of the two-body problem should be embedded into the Mean Field realm. Therefore the scattering length a should be understood as the quantity renormalized by the medium. How to perform such a renormalization is a separate problem not discussed here.

The BEC and BCS domains are separated by the “unitary point” $(k_F a)^{-1} = 0$. In the vicinity of this point the scattering length a characterizes the size of the pair. Our next task is to demonstrate how the “connection problem” is solved, i.e., describe the BEC-BCS crossover.

4 BCS-BEC Crossover

The continuous evolution from BCS to BEC regime is called the BCS-BEC crossover. Such a transition takes place either by increasing the strength of the interaction or by decreasing the carrier density. The fact that the BCS wave function may undergo a smooth evolution and describe the tightly bound fermion pairs was first noticed long ago [42]-[46]. According to [46], the remark that “there exists a transformation that carries the BCS into BE state” was originally made by F.J.Dyson in 1957 (i.e., the same year that the BCS paper [22] was published). The BCS-BEC crossover for quarks was first discussed in [7].

The explicit transformation from the BCS solution (14) (weak coupling) to the Schrodinger equation for molecular-like state (strong coupling) was performed in [44]. Here we outline a relativistic generalization of this procedure suited for light quarks [33]. We omit the details of the derivation since it is similar to the derivation of the propagator described in textbooks. One should take the derivative over d^* of $\ln Z_B$ (see (15)). Then at $T = 0$ the positive and negative frequencies relativistic wave functions in momentum space have the form

$$\varphi(k) = \frac{d(k)}{\varepsilon_+(k)}, \quad \chi(k) = \frac{d(k)}{\varepsilon_-(k)}, \quad (23)$$

where the quantities $\varepsilon_{\pm}(k)$ are defined after Eq.(16). The momentum depen-

dence of $d(k)$ means that we have assumed that the four-fermion interaction (see (3)) is not point-like any more. Then we use the self-consistency relation (16) and obtain the following set of coupled equations for φ and χ

$$(\sqrt{k^2 + m^2} - \mu)\varphi(k) = (1 - 2n_k) \int \frac{d\mathbf{p}}{(2\pi)^3} g(p - k)[\varphi(p) + \chi(p)], \quad (24)$$

$$(\sqrt{k^2 + m^2} + \mu)\chi(k) = (1 - 2\bar{n}_k) \int \frac{d\mathbf{p}}{(2\pi)^3} g(p - k)[\chi(p) + \varphi(p)], \quad (25)$$

$$1 - 2n_k = \frac{\sqrt{k^2 + m^2} - \mu}{\varepsilon_+(k)}, \quad 1 - 2\bar{n}_k = \frac{\sqrt{k^2 + m^2} + \mu}{\varepsilon_-(k)}. \quad (26)$$

To obtain the Schrodinger equation we neglect the negative frequency component in (24), expand the square root and consider the dilute limit. The result is

$$\left(\frac{k^2}{2m} - \tilde{\mu}\right)\varphi(k) = \int \frac{d\mathbf{p}}{(2\pi)^3} g(p - k)\varphi(p), \quad (27)$$

where $\tilde{\mu} = \mu - m$ in line with the definition of the chemical potential in nonrelativistic case.

In relativistic case the phase diagram in the $(n_k/\bar{n}_k, \mu)$ plane has two symmetric branches corresponding to quarks and antiquarks with possible “exciton-like” instability.

In order to keep these two lectures down to a reasonable length we have not discussed the physics of ultra cold fermionic atoms at length. This subject is rapidly developing and one should follow the current literature. The last remark we wish to make concerns Eq. (22) and Efimov effect [47]. The point is that Feshbach resonance may give rise to infinite number (proportional to $\ln(|a|/r_0)$) of three body atomic states.

These lectures were delivered at 8th Moscow School of Physics in early spring of 2005. The author wishes to express his deep gratitude to the Organizing Committee for the invitation and a nice time spent there.

References

- [1] M.Alford, K.Rajagopal and F.Wilczek, Phys. Lett. **B 422**, 247 (1998).

- [2] R.Rapp, T.Schafer, E.V.Shuryak and M.Velkovsky, Phys. Rev. Lett. **81**, 53 (1998).
- [3] M.Greiner, C.A.Regal, D.S.Jin, Nature, **426**, 537 (2003).
- [4] S.Jochim et al., Science, **302**, 2101 (2003).
- [5] M.W.Zwierlein et al., Phys. Rev. Lett. **91**, 250401 (2003).
- [6] B.Kerbikov, arXiv:hep-ph/0110197.
- [7] B.Kerbikov, Phys. Atom. Nucl. **65**, 1918 (2002).
- [8] K.Rajagopal and F.Wilczek, In: At the Frontier of Particle Physics. Handbook of QCD, Ed. by M.Shifman, World Sci., Singapore, 2001, Vol. 3, p.2061;
M.Alford, Annu. Rev. Nucl. Part. Sci. **51**, 131 (2001);
G.Nardulli, Riv.Nuovo Cim. **25**, 1 (2002);
I.A.Shovkovy, ArXiv:nucl-th/0410091.
- [9] Qijin Chen et.al., Phys. Repts. **412**, 1 (2005).
- [10] M.H.Anderson et al., Science, **269**, 198(1995).
- [11] C.C.Bradley et al., Phys. Rev. Lett. **75**, 1687 (1995).
- [12] K.B.Davis et al., Phys. Rev. Lett. **75**, 3969 (1995).
- [13] F.Dalfovo, S.Giorgini, L.P.Pitaevsky, and S.Stringari, Rev. Mod. Phys. **71**, 463 (1999).
- [14] G.Baym, Nucl. Phys. **A751**, 373 (2005).
- [15] L.D.Carr, G.V.Shlyapnikov and Y.Castin, Phys. Rev. Lett. **92**, 150404 (2004).
- [16] Qijin Chen, Jelena Stajic, and K.Levin, ArXiv: cond.-mat/0411090.
- [17] M.R.Schafroth, Phys. Rev. **96**, 1442 (1954).
- [18] J.Polchinsky, ArXiv: hep-th/9210046 (Lectures presented at TASI 1992).

- [19] S.Weinberg, Nucl. Phys. **B 413**, 567 (1994).
- [20] A.Svidzinsky, Theor. Matem. Fiz. **9**, 273 (1971).
- [21] C.N. Yang, Rev. Mod. Phys. **34** , 694 (1962).
- [22] J.Bardeen, L.N.Cooper, J.R.Schrieffer, Phys. Rev. **108**, 1175 (1957).
- [23] C.P.Enz, A Course on Many-Body Theory Applied to Solid-State Physics, World Scientific, Singapore, 1992.
- [24] P.W.Anderson, Rev. Mod. Phys. **38**, 298 (1966).
- [25] J.I.Kapusta, Finite Temperature Field Theory, Cambridge University Press, 1989.
- [26] V.N.Popov, Functional Integrals in Quantum Field theory and Statistical Physics, Reidel, Doordrecht, 1983.
- [27] R.Haussmann, Z.Phys. **B91**, 291 (1993).
- [28] J.Berges and K.Rajagopal, Nucl. Phys. **B538**, 215 (1999).
- [29] B.Kerbikov, arXiv:hep-ph/0106324.
- [30] A.V.Svidzinsky, Nonuniform Problems of the Superconductivity Theory, Nauka, Moscow, 1982.
- [31] V.M.Loktev, R.M.Quick, S.G.Sharapov, Phys. Repts. **349**, 1 (2001).
- [32] F. Pistolesi and G.C.Strinati, Phys. Rev. **B 53**, 15168 (1996).
- [33] B.Kerbikov, Phys. Atom. Nucl. **68**, 890 (2005).
- [34] A.Larkin and A.Varlamov, in: The Physics of Conventional and Unconventional Superconductors. Eds. K.H.Bennermann, J.B.Ketterson, Springer-Verlay, Berlin-Heidelberg, 2001.
- [35] N.Agasian, B.Kerbikov, V.Shevchenko, Phys. Repts. **320**, 131 (1999).
- [36] Yu.A.Simonov, Phys. Usp. **39**, 313 (1997).
- [37] B.I.Halperin, T.C.Lubensky, Shang-keng Ma, Phys. Rev. Lett. **32**, 292 919740.

- [38] B.Kerbikov, Talk at INT Program 04-1,
<http://www.int.washington.edu/talks>.
- [39] T.Matsuura, K.Iida, T.Hatsuda, G.Baym, Phys. Rev. **D69**, 074012 (2004).
- [40] M.Holland et al., Phys. Rev. Lett. **87**, 120406 (2001).
- [41] E.Timmermans et al., Phys. Lett. **A285**, 228 (2001).
- [42] D.M.Eagles, Phys. Rev. **186**, 456 (1969).
- [43] A.J.Leggett in: Modern Trends in the Theory of condensed matter, Springer-Verlag, Berlin, 1980.
- [44] P.Nozieres and S.Schmitt-Rink, J.Low, Temp. Phys. **59**, 195 (1985).
- [45] P.W.Anderson and W.F.Brinkman in: Proceedings of the 15th Scottish University summer school in Physics, Acad. Press, New York, 1975.
- [46] P.W.Anderson, Basic Notions of Condensed Matter physics, Benjamin/Cummings Publ., Menlo Park, 1983, p.105.
- [47] V.Efimov, Phys. Lett. **33B**, 563 (1970).